

On Group-Theoretic Finite-Mode Approximation of $2D$ Ideal Hydrodynamics

Zbigniew Peradzyński¹ Hanna E. Makaruk²
Robert M. Owczarek^{2*}

¹ University of Warsaw, Department of Mathematics, Banacha 2, 02-097 Warsaw, Poland, and Institute of Fundamental Technological Research, Polish Academy of Sciences, Świętokrzyska 21, 00-041 Warsaw, Poland

² E-ET, MS M319, Los Alamos National Laboratory, Los Alamos, NM 87545, USA, on leave from Institute of Fundamental Technological Research, Polish Academy of Sciences, Świętokrzyska 21, 00-041 Warsaw, Poland

Abstract

Structure constants of the $su(N)$ (N odd) Lie algebras converge when N goes to infinity to the structure constants of the Lie algebra $sdiff(T^2)$ of the group of area-preserving diffeomorphisms of a $2D$ torus. Thus Zeitlin and others hypothesized that solutions of the Euler equations associated with $su(N)$ algebras converge to solutions of the Euler equations of incompressible fluid dynamics on a $2D$ torus. In the paper we prove the hypothesis. Our numerical experiments show the Galerkin method applied to Euler equation of hydrodynamics is computationally more efficient in the range of time in which it is stable than that based on the $SU(N)$ approximation. However, the latter is stable for much longer time. These numerical results agree with theoretical expectations.

PACS Numbers:

03.40.60 Gc— Fluid dynamics;

*corresponding author: rmo@lanl.gov, Los Alamos NM 87545, USA

02.70.Dh — Finite element and Galerkin methods

02.20.a — Group theory

02.20.Tw — Infinite dimensional Lie groups.

1 Introduction

In the beginning of 90's a finite-mode approximation of $2D$ ideal hydrodynamics was introduced ([1], [2], [3]). The approximation is based on convergence of the structure constants of the Lie algebras $su(N)$ (N -odd), when $N \rightarrow \infty$ to the structure constants of the Lie algebra $sdiff(T^2)$. The Lie algebra $sdiff(T^2)$ consists of divergenceless nonconstant vector fields on a $2D$ torus. The vector fields are infinitesimal generators of the group of area-preserving diffeomorphisms of the $2D$ torus. Incidentally, the group can be also identified as the group of symplectic diffeomorphisms, for the natural symplectic structure of the torus. The group is the basic symmetry group for incompressible ideal hydrodynamics on a $2D$ torus. In the physical model the fluid is maintained in a region of a plane, appropriate boundary conditions being satisfied. The approximation of $sdiff(T^2)$ by $su(N)$ is used quite widely both in hydrodynamics and in string theory [1, 2, 3, 4, 5, 6, 7, 8]. There are also other finite-mode approximations of the Euler and similar equations related to symplectic geometry [9, 10, 11, 12].

In [13] the approach to approximate $sdiff(T^2)$ is formalized and a weak uniqueness theorem is proved. Unfortunately, the weak uniqueness does not lead to an isomorphism of the limit algebras. Indeed, there are examples of various limit algebras which have their structure constants equal to the structure constants of some infinite dimensional algebras and despite this equality the limit algebras are not isomorphic to the infinite dimensional ones. As shown in [13], the $\lim_{N \rightarrow \infty} su(N)$ is neither isomorphic to $sdiff(T^2)$ nor to $sdiff(S^2)$, where $sdiff(S^2)$ consists of divergenceless vector fields on the

2D sphere S^2 . This approximation is still not well understood and additional work should be done towards its satisfactory formulation.

The plan of the paper is as follows.

In the second section we discuss the issues connected with approximating the Lie algebra $sdiff(T^2)$ by the Lie algebras $su(N)$. We point out weaknesses of this approximation.

In the third section we remind how the Euler equations are being associated with arbitrary Lie algebras. Then we specify the Lie algebras to be $su(N)$ (N odd) and $sdiff(T^2)$. Despite the weak form of convergence of $su(N)$ to $sdiff(T^2)$, solutions of the Euler equations associated to the Lie algebras $su(N)$ converge, in a sense defined further in the paper, when N goes to infinity, to the solutions of the Euler equation associated with the Lie algebra $sdiff(T^2)$. Proof of this fact is the central result of this paper.

In the fourth section we discuss numerical calculations within this approach and within the Galerkin approximation. Numerical stability of this method in comparison to Galerkin one turns out to be significantly better. On the other hand, the numerical simulations demand significantly more calculation time than the Galerkin one.

In the Conclusions we discuss the finite-mode approximation based on the group theoretic relations as a method of choice for solving Euler equation whenever numerical calculations are expected to give reliable results for long evolution times, and adequate computational power is available.

We supplement the paper with an Appendix, in which we discuss geometry of noncommutative tori, providing geometric interpretation of the approximation method.

2 Approximation of the algebra $sdiff(T^2)$ by $su(N)$ algebras

In this section we remind the formal approach to the approximation of $sdiff(T^2)$ and $sdiff(S^2)$ algebras given in [13] and called there L_α -convergence. We remind here the result on weak uniqueness (proposition 3.3 from [13]). We also discuss behavior of generators of $su(N)$ in the form chosen in [1] when N goes to infinity. We show that their norms diverge in this limit.

In [13] instead of algebras $u(N)$, $su(N)$ are considered their complexifications, $gl(N, \mathcal{C})$ and $sl(N, \mathcal{C})$. It is quite a standard to consider instead of algebras $sdiff(T^2)$, $sdiff(S^2)$, etc., their complexifications, so the above procedure is fully justified. It does not lead to any principal differences in comparison with our dealing with corresponding real algebras, since the structure constants remain the same and the relations of being isomorphic or non-isomorphic are preserved by complexification.

Let us begin with giving necessary definitions of objects appearing in this section. First, let us remind some definitions and basic facts about the Lie algebras $gl(\infty)$, $gl_+(\infty)$, and L_Λ , as they are presented in [13].

First, $gl(\infty)$ is defined as $gl(\infty) := \{(a_{ij})_{ij \in \mathbb{Z}} : a_{ij} \in \mathcal{C}, \text{ all but finite number of } a_{ij} = 0\}$. This is then a matrix algebra with finite support. As usual for matrix Lie algebras the commutator of matrices serves as the Lie bracket.

The Lie algebra $gl_+(\infty)$ is defined in the same way with the only difference that the indices of the elements of the matrices in the Lie algebra are not integers $\mathbb{Z}$ but natural numbers $\mathbb{N}$. Since $\mathbb{Z}$ and $\mathbb{N}$ are bijective, any such bijection induces an isomorphism of $gl(\infty)$ and $gl_+(\infty)$. However, the isomorphisms are not canonical, which means they depend on the choice of a basis in each of the Lie algebras. Since $gl(\infty)$ and $gl_+(\infty)$ have only finite number of elements different from zero, there is no problem with defining the

trace of the elements in these Lie algebras. Namely, the number of non-zero diagonal elements is also finite for all elements of $gl(\infty)$ and $gl_+(\infty)$, and because of that the standard definition of a trace as the sum of diagonal elements of a matrix is unchanged for both algebras. Then the Lie algebras $sl(\infty)$ and $sl_+(\infty)$ can be easily defined as consisting of those elements of $gl(\infty)$, $gl_+(\infty)$, which have trace equal to zero, and with the Lie structure inherited from the algebras $gl(\infty)$, $gl_+(\infty)$, respectively.

The Lie algebras $sdiff(T^2)$ and L_Λ , which are important in our further considerations, are defined in the following way. One begins with the complex vector space:

$$V = \{T_{\bar{m}}, \bar{m} \in \mathbb{Z} \times \mathbb{Z}\}, \quad (1)$$

it means a complex vector space generated by the basis $T_{\bar{m}}$, with indices $\bar{m} = (m_1, m_2)$ such that m_1 and m_2 are integers.

The following interesting Lie algebra structures, indexed by $\Lambda \in \mathbb{R}$, can be defined on V : For $\Lambda \in \mathbb{R} - \{0\}$ they are defined on the generators $T_{\bar{m}}$ by:

$$[T_{\bar{m}}, T_{\bar{n}}]^\Lambda := \frac{1}{2\pi\Lambda} \sin(2\pi\Lambda(\bar{m} \times \bar{n})) T_{\bar{m}+\bar{n}} \quad (2)$$

and for $\Lambda = 0$ by:

$$[T_{\bar{m}}, T_{\bar{n}}]^0 := (\bar{m} \times \bar{n}) T_{\bar{m}+\bar{n}}, \quad (3)$$

where

$$\bar{m} \times \bar{n} = m_1 n_2 - m_2 n_1 \quad \text{for} \quad \bar{m} = (m_1, m_2), \quad \bar{n} = (n_1, n_2). \quad (4)$$

Then, as usual, the Lie brackets are extended to the whole V by linearity. These Lie algebras are denoted $\tilde{L}_\Lambda = (V, [\cdot, \cdot]^\Lambda)$. They are direct sums of subalgebras

$$\tilde{L}_\Lambda = \{T_{(0,0)}\} \oplus \{T_{\bar{m}} : \bar{m} \in \mathbb{Z} \times \mathbb{Z} - \{0\}\} \quad (5)$$

where $\{\cdot\}$ denotes a vector space spanned on generators listed inside the figure brackets, and the Lie structure is defined by restriction of $[\cdot, \cdot]^\Lambda$ to the given subspace of V .

The Lie algebra L_Λ is defined as the second summand in the above sum, $L_\Lambda = (\{T_{\bar{m}} : \bar{m} \in \mathbb{Z} \times \mathbb{Z} - \{0\}\}, [\cdot, \cdot]^\Lambda)$. In the case $\Lambda = 0$, L_o is identified with the complexified Lie algebra of the nonconstant divergenceless vector fields on T^2 , which is the $sdiff(T^2)$.

Since approximation of the Lie algebras L_Λ , $sdiff$, etc. by the algebras $gl(n)$ or $su(n)$ can not be realized in a strong sense of inductive or projective limit as it is often done for other infinite-dimensional algebras, the authors of [13] axiomatized the concept of so called L_α -limit of the Lie algebras. This notion is a generalization of intuitively defined limits from the papers [8], [14].

The definition consists of two axioms. The first axiom demands existence of a special family of real or complex Lie algebras $(\mathcal{G}_\alpha, [\cdot, \cdot]_\alpha, \alpha \in I)$, where the index set $I = \mathbb{N}$ or $\mathbb{R}$. This family should be equipped with a family of metrics d_α . Accordingly to the axiom there should exist another real or complex Lie algebra $(\mathcal{G}, [\cdot, \cdot])$ such that for every $\alpha \in I$ there exists a surjective map $p_\alpha : \mathcal{G} \mapsto \mathcal{G}_\alpha$ and the condition

$$\forall x, y \in \mathcal{G} \text{ if } d_\alpha(p_\alpha(x), p_\alpha(y)) \xrightarrow{\alpha \rightarrow \infty} 0, \text{ then } x = y. \quad (6)$$

should be satisfied. The second axiom concerns the definition of a Lie algebra structure on the L_α -limit. Accordingly to this axiom $(\mathcal{G}, [\cdot, \cdot])$, is called L_α -limit of the algebras $(\mathcal{G}_\alpha, [\cdot, \cdot]_\alpha, d_\alpha, \alpha \in I)$ and the family of the algebras is called the approximating sequence introduced by $(p_\alpha, \alpha \in I)$ if the condition

$$d_\alpha(p_\alpha[x, y], [p_\alpha x, p_\alpha y]_\alpha) \xrightarrow{\alpha \rightarrow \infty} 0. \quad (7)$$

is satisfied. There (in [13]) was also proved a weak uniqueness theorem, which states that if $(\mathcal{G}_\alpha, [\cdot, \cdot]_\alpha, d_\alpha, \alpha \in I)$ is an approximating sequence for $(\mathcal{G}, [\cdot, \cdot])$ induced by $(p_\alpha, \alpha \in I)$, and $\mathcal{G}'$ is a linear subspace of $\mathcal{G}$, which carries a Lie structure $[\cdot, \cdot]'$ which projects on each $\mathcal{G}_\alpha$, then $(\mathcal{G}', [\cdot, \cdot]')$ is a Lie subalgebra

of $(\mathcal{G}, [\cdot, \cdot])$ if and only if the approximating sequence for $\mathcal{G}$ is by restriction also approximating sequence for $\mathcal{G}'$ induced by the restriction of the p_α .

However, this theorem about weak uniqueness is not of very big value, because it does not guarantee that if we start with a given approximating sequence of the Lie algebras $(\mathcal{G}_\alpha, \alpha \in I)$ we end up with isomorphic limit algebras, independently of how we came to the limits.

Let us remind how $sdiff(T^2)$ is described rigorously within the L_α limit of the algebras L_Λ when $\Lambda \rightarrow 0$. One begins with the sequence of algebras $L^N = \tilde{L}_{1/N}$, i.e. the sequence of algebras $\tilde{L}_\Lambda$ with $\Lambda = \frac{1}{N}$. One distinguishes an ideal J^N in each algebra L^N of the sequence:

$$J^N := \{T_{\bar{m}} - T_{\bar{m}+N\bar{a}} : \bar{m}, \bar{a} \in \mathbb{Z} \times \mathbb{Z}\}. \quad (8)$$

Then one defines the sequence of factor algebras $L_{(N)} := L^N / J^N$ with $\pi_N : L^N \rightarrow L_{(N)}$ the canonical projection. As the basis of each algebra in the sequence serve $\pi_N(T_{\bar{m}})$, $\bar{m} = (m_1, m_2)$, $0 \leq m_1, m_2 < N$. The Lie structure in each algebra of the sequence is given by:

$$[\pi_N(T_{\bar{m}}), \pi_N(T_{\bar{n}})]^{(N)} = \frac{N}{2\pi} \sin\left(\frac{2\pi}{N}(\bar{m} \times \bar{n})\right) \pi_N(T_{(\bar{m}+\bar{n}) \bmod N}). \quad (9)$$

This algebra $L_{(N)}$ is exactly identical with the algebra $gl(N)$ with N odd. The approximating sequence consists then of the algebras $(L_{(N)}, [\cdot, \cdot]^{(N)})$ with the index set $I = \mathbb{N}$. The surjective maps p_α are identified in this case as the maps π_N . The metric on each $L_{(N)}$ is induced by defining it on the generators by

$$\langle \pi_N(T_{\bar{m}}), \pi_N(T_{\bar{n}}) \rangle = \delta_{m_1, n_1} \delta_{m_2, n_2}. \quad (10)$$

and then extending to the whole $L_{(N)}$ by linearity.

It is shown in [13] that the 2 axioms of L_α -limit are satisfied for this sequence. Therefore, the algebra $sdiff(T^2)$ is an L_α -limit of the algebras $gl(N, \mathcal{C})$. Nevertheless, $gl(\infty)$ which is an inductive limit of the algebras

$gl(N, \mathbb{C})$, so it is the limit in a strong sense, is not isomorphic to $sdiff(T^2)$. This fact is also proved in [13].

The above example shows that one should be careful with approximating infinite-dimensional real algebras by finite dimensional ones in such weak sense as the L_α -limit. We will demonstrate now another difficulty of this definition of a limit. The difficulty comes from the fact that the renormalized basis of the Lie algebra $su(N)$ has divergent norm for N going to infinity. To be more precise, let us remind that the basis of the $su(N)$ algebra which leads to the mentioned L_α -approximation of $sdiff(T^2)$ is defined as:

$$\mathcal{J}_{\bar{n}} = \lambda^{n_1 n_2 / 2} g^{n_1} h^{n_2}, \quad (11)$$

where $\lambda = e^{4i\pi/N}$, $g = \text{diag}(1, \lambda, \dots, \lambda^{N-1})$, and

$$h = \begin{bmatrix} 0 & 1 & . & . & . & 0 \\ 0 & 0 & 1 & . & . & 0 \\ . & . & . & . & . & . \\ 1 & . & . & . & . & 0 \end{bmatrix}$$

is a cyclic permutation matrix. Then, to obtain desired form of the structure constants of the algebra, a renormalization is done by multiplying all the basis elements by $\frac{iN}{4\pi}$. As a result, the renormalized basis elements for $N \rightarrow \infty$ become elements of infinite norm. This is in apparent contrast to the behavior of the basis elements of the algebra $sdiff(T^2)$

$$L_{\bar{n}} = i \exp(i\bar{n} \cdot \bar{x})(n_2 \partial_1 - n_1 \partial_2) \quad (12)$$

, which are all of finite norm.

It seems then that the algebraic arguments on approximation of $sdiff(T^2)$ by $su(N)$ need careful treatment.

What is also of interest, especially in the context of geometric description of an ideal incompressible fluid, and its geometric quantization like in [15] or [16], it is a better understanding of convergence of coadjoint orbits of the

groups $SU(N)$ to the coadjoint orbits of the group $SDiff(T^2)$. Analogous interesting question concerns the Casimir functions. These in the case of $SU(N)$ are: $tr A^2, tr A^3, \dots, tr A^n, \dots$ for an $A \in su(N)$. Their counterparts for the $SDiff(T^2)$ are: $\int_{T^2} \omega^2 \mu, \int_{T^2} \omega^3 \mu, \dots, \int_{T^2} \omega^n \mu, \dots$, where μ is the standard surface element of the torus T^2 . The work on this issue is in progress.

3 Approximating Euler equation

In this section we introduce the Euler equation for incompressible flows of an ideal fluid on a $2D$ torus. Then we remind the definition of the Euler equations connected with arbitrary Lie algebras, due to Arnold. Next, we specify the latter equations to the case of $su(N)$ Lie algebras (N odd) and to the case of the Lie algebra $sdiff(T^2)$. Finally, we prove the theorem on convergence of solutions of the Euler equations for $su(N)$ to the solutions of the Euler equation connected with the $sdiff(T^2)$ Lie algebra.

As was noticed in [17], the Euler equations can be generalized to any Lie algebra by

$$\dot{\omega}_i = g^{jk} C_{ik}^r \omega_j \omega_r, \quad (13)$$

where C_{ik}^r are structure constants of the algebra and g is a symmetric metric. For example, g is the inverse of the inertia tensor in the case of motion of a rigid body, for which the appropriate algebra is $so(3)$. The summation over repeating indices is understood.

The equation for two-dimensional hydrodynamics of an ideal fluid can be expressed in terms of the Poisson bracket $\{\cdot, \cdot\}$ of vorticity ω and the stream function as follows

$$\dot{\omega} = \{\Delta^{-1} \omega, \omega\}, \quad (14)$$

where $\Delta^{-1} \omega$ denotes the stream function, i.e. the solution of $\Delta \psi = \omega$. In the case of the flow in a rectangle with periodic boundary condition (i.e. on

a two-dimensional torus T^2) we assume the mean vorticity equal to zero, $\int_{T^2} \omega = 0$. As noticed by Zeitlin, the symplectic diffeomorphisms of T^2 are generated by

$$L_{\bar{n}} = i \exp[i(n_1 x_1 + n_2 x_2)](n_2 \partial_1 - n_1 \partial_2) = -i \exp(i\bar{n} \cdot \bar{x})(\bar{n} \times \nabla),$$

where $\bar{n}^2 = (n_1^2 + n_2^2) \neq 0$. These operators satisfy the Jacobi identity and form the Lie algebra $sdiff(T^2)$. For the commutators we have

$$[L_{\bar{n}}, L_{\bar{m}}] = (\bar{n} \times \bar{m}) L_{\bar{n}+\bar{m}}, \quad \text{where} \quad \bar{n} \times \bar{m} = n_1 m_2 - n_2 m_1. \quad (15)$$

Equation (14), when expressed in the basis (12), i.e. in terms of the Fourier components, $\{\omega_{\bar{n}}\}$ of ω , takes the form

$$\dot{\omega}_{\bar{m}} = \sum_{\bar{k}} \frac{\bar{m} \times \bar{k}}{k^2} \omega_{\bar{m}+\bar{k}} \omega_{-\bar{k}}, \quad (16)$$

where $\omega_{-\bar{k}} = \omega_{\bar{k}}^*$, because $\omega(\bar{x})$ is real. Taking (11) as the basis for the Lie algebra $su(N)$ for $N = 2n + 1$ one can construct a system which formally approximates the system (16). Assuming $L_{\bar{n}} = \frac{iN}{4\pi} \mathcal{J}_{\bar{n}}$, where $\mathcal{J}_{\bar{n}}$ are defined in (11) we have

$$[L_{\bar{n}}, L_{\bar{m}}] = \frac{N}{2\pi} \sin\left(\frac{2\pi}{N}\right)(\bar{n} \times \bar{m}) L_{\bar{n}+\bar{m} \bmod N} \quad (17)$$

in which the summation $\bmod N$ in both components is understood. Thus, assuming the inertia tensor analogous to the one in (16) we may define the following Euler equations on the dual of $su(N)$

$$\dot{\omega}_{\bar{m}} = \sum_{\bar{k}} \frac{\sin[2\pi(\bar{m} \times \bar{k})/N]}{\frac{2\pi}{N} k^2} \omega_{\bar{m}+\bar{k} \bmod N} \omega_{-\bar{k}}, \quad (18)$$

where $-\frac{N-1}{2} \leq k_i, m_i \leq \frac{N-1}{2}$, $i = 1, 2$. In the literature the system (18) is known as a finite dimensional analog of two-dimensional hydrodynamics. However, no proof was known that solutions of (18) are indeed approximating those of (16). To prove that such a correspondence exists we will treat

solutions of (18) as infinite sequences assuming $\omega_{\bar{k}}, \omega_{-\bar{k}}$ are zero for $|k_i| > \frac{N-1}{2}$. Let us assume therefore that $\{\omega_{\bar{k}}(t)\}$ is a solution of Eqs(16) with the initial data $\{\omega_{\bar{k}}(0)\}$. Correspondingly, let $\{\tilde{\omega}_{\bar{k}}(t)\}$ be a solution of Eqs(18) with initial values $\{\tilde{\omega}_{\bar{k}}(0)\}$. We will assume that the difference $\omega_{\bar{k}}(0) - \tilde{\omega}_{\bar{k}}(0)$ is small in l^2 i.e. $\sum_{\bar{k}} |\omega_{\bar{k}}(0) - \tilde{\omega}_{\bar{k}}(0)|^2 < \delta$. Then we will prove under some additional assumptions for ω that on a finite time interval the solutions are also close. Let us write the equation for the difference $\Omega(t) = \omega(t) - \tilde{\omega}(t)$. Subtracting (18) from (16) one obtains after some manipulations

$$\begin{aligned} \dot{\Omega}_{\bar{m}} = & \sum_{\bar{k}} \frac{\bar{m} \times \bar{k}}{\bar{k}^2} \left(1 - \frac{\sin(\varepsilon \bar{m} \times \bar{k})}{\varepsilon \bar{m} \times \bar{k}} \right) \omega_{\bar{m}+\bar{k}} \omega_{-\bar{k}} + \\ & \sum_{\bar{k}} \frac{\sin(\varepsilon \bar{m} \times \bar{k})}{\varepsilon \bar{k}^2} \{ \Omega_{\bar{m}+\bar{k}} \omega_{-\bar{k}} + \tilde{\omega}_{\bar{m}+\bar{k}} \Omega_{-\bar{k}} \}, \end{aligned} \quad (19)$$

where $\varepsilon = \frac{2\pi}{N}$. It is desirable to express the r.h.s. of (19) only in terms of variables $\Omega_{\bar{k}}, \omega_{\bar{k}}$. Since $\omega_{\bar{m}+\bar{k}} = \tilde{\omega}_{\bar{m}+\bar{k}} + \Omega_{\bar{m}+\bar{k}}$, we have

$$\begin{aligned} \dot{\Omega}_{\bar{m}} = & \sum_{\bar{k}} \frac{\bar{m} \times \bar{k}}{\bar{k}^2} \left(1 - \frac{\sin(\varepsilon \bar{m} \times \bar{k})}{\varepsilon \bar{m} \times \bar{k}} \right) \omega_{\bar{m}+\bar{k}} \omega_{-\bar{k}} + \\ & \sum_{\bar{k}} \frac{\sin(\varepsilon \bar{m} \times \bar{k})}{\varepsilon \bar{k}^2} \{ \Omega_{\bar{m}+\bar{k}} \omega_{-\bar{k}} + \omega_{\bar{m}+\bar{k}} \Omega_{-\bar{k}} - \Omega_{\bar{m}+\bar{k}} \Omega_{-\bar{k}} \}, \end{aligned} \quad (20)$$

Now we are ready to derive the equation for the L^2 -norm of Ω

$$\frac{1}{2} \|\Omega\|^2 = \frac{1}{2} \sum_{\bar{m}} |\Omega_{\bar{m}}|^2 = \frac{1}{2} \sum_{\bar{m}} \Omega_{\bar{m}} \Omega_{-\bar{m}}. \quad (21)$$

The last equality in the chains follows from $\Omega(\bar{x})$ being real.

Let us observe that

$$\frac{1}{2} \frac{d}{dt} \sum_{\bar{m}} \Omega_{\bar{m}} \Omega_{-\bar{m}} = \frac{1}{2} \sum_{\bar{m}} \left(\frac{d\Omega_{\bar{m}}}{dt} \Omega_{-\bar{m}} + \Omega_{\bar{m}} \frac{d\Omega_{-\bar{m}}}{dt} \right) = \sum_{\bar{m}} \Omega_{\bar{m}} \frac{d\Omega_{-\bar{m}}}{dt} \quad (22)$$

,since the sum $\sum_{\bar{m}} \Omega_{\bar{m}} \frac{d\Omega_{-\bar{m}}}{dt}$ by substitution $\bar{m} \rightarrow -\bar{m}$ becomes $\sum_{\bar{m}} \Omega_{-\bar{m}} \frac{d\Omega_{\bar{m}}}{dt}$.

Multiplying Eq.(20) by $\Omega_{-\bar{m}}$, performing the summation over $\bar{m}$ one obtains

$$\frac{1}{2} \frac{d}{dt} \|\Omega(t)\|^2 = \sum_{\bar{m}} a_{\bar{m}}(\varepsilon) \Omega_{-\bar{m}}$$

$$\begin{aligned}
& + \sum_{\bar{m}} \sum_{\bar{k}} \frac{\sin(\varepsilon \bar{m} \times \bar{k})}{\varepsilon \bar{k}^2} \omega_{\bar{m}+\bar{k}} \Omega_{-\bar{k}} \Omega_{-\bar{m}} \\
& + \sum_{\bar{m}} \sum_{\bar{k}} \frac{\sin(\varepsilon \bar{m} \times \bar{k})}{\varepsilon \bar{k}^2} \Omega_{\bar{m}+\bar{k}} \Omega_{-\bar{m}} \omega_{-\bar{k}} \\
& - \sum_{\bar{m}} \sum_{\bar{k}} \frac{\sin(\varepsilon \bar{m} \times \bar{k})}{\varepsilon \bar{k}^2} \Omega_{\bar{m}+\bar{k}} \Omega_{-\bar{m}} \Omega_{-\bar{k}}
\end{aligned} \tag{23}$$

where

$$a_{\bar{m}}(\varepsilon) = \sum_{\bar{k}} a_{\bar{m}}^{\bar{k}}(\varepsilon) \omega_{\bar{m}+\bar{k}} \omega_{-\bar{k}}. \tag{24}$$

and where

$$a_{\bar{m}}^{\bar{k}}(\varepsilon) = \frac{\bar{m} \times \bar{k}}{\bar{k}^2} \left(1 - \frac{\sin(\varepsilon \bar{m} \times \bar{k})}{\varepsilon \bar{m} \times \bar{k}} \right). \tag{25}$$

In the last two summands repeats the term:

$$I_{\bar{k}} := \sum_{\bar{m}} \sin(\varepsilon \bar{m} \times \bar{k}) \Omega_{\bar{m}+\bar{k}} \Omega_{-\bar{m}} \tag{26}$$

Changing the name of the summation index does not change the value of $I_{\bar{k}}$. Let us then introduce the change from $\bar{m}$ into $-(\bar{m} + \bar{k})$. As a result one gets $(\bar{m}' = -(\bar{m} + \bar{k}))$ and then we omit $'$.

$$\begin{aligned}
I_{\bar{k}} &= \sum_{\bar{m}} \sin(\varepsilon (-(\bar{m} + \bar{k}) \times \bar{k})) \Omega_{-\bar{m}} \Omega_{\bar{m}+\bar{k}} \\
&= - \sum_{\bar{m}} \sin(\varepsilon (\bar{m} \times \bar{k})) \Omega_{\bar{m}+\bar{k}} \Omega_{-\bar{m}} = -I_{\bar{k}}
\end{aligned} \tag{27}$$

Therefore $I_{\bar{k}} \equiv 0$.

Now we will prove that the sequence $\{a_{\bar{m}}(\varepsilon)\}$ defined by (24) satisfies $\sum |a_{\bar{m}}|^2 = \delta(\varepsilon) \rightarrow 0$ when $\varepsilon \rightarrow 0$ provided that the solution $\omega(x)$ belongs to the Sobolev space¹ $H^\sigma(T^2)$ for $\sigma > 2$. We have in such a case $\sum_{\bar{m}} |\omega_{\bar{m}}|^2 (1 + \bar{m}^2)^\sigma < \infty$ and therefore $\omega_{\bar{m}}$ can be represented as $\omega_{\bar{m}} = \frac{c_{\bar{m}}}{(1 + \bar{m}^2)^{\sigma/2}}$ where still $\sum_{\bar{k}} |c_{\bar{k}}|^2 = c < \infty$. Thus denoting $\beta_{\bar{m}}^{\bar{k}}(\varepsilon) = a_{\bar{m}}^{\bar{k}}(\varepsilon) (1 + (\bar{m} + \bar{k})^2)^{-\sigma/2} (1 +$

¹We know [18] that in two-dimensional case the Euler equations of an ideal fluid have global solutions whose quality (differentiability properties) are related to the quality of initial data.

$\bar{k}^2)^{-\sigma/2}$ we have $a_{\bar{m}} = \sum_{\bar{k}} \beta_{\bar{m}}^{\bar{k}}(\varepsilon) c_{\bar{m}+\bar{k}} c_{-\bar{k}}$, which by Schwartz inequality gives us

$$|a_{\bar{m}}| \leq \sqrt{\sum_{\bar{k}} |\beta_{\bar{m}}^{\bar{k}}(\varepsilon)|^2} \sqrt{\sum_{\bar{k}} |c_{\bar{m}+\bar{k}} c_{-\bar{k}}|^2}. \quad (28)$$

The last sum on the right hand side is obviously bounded because

$$\sum_{\bar{k}} |c_{\bar{m}+\bar{k}} c_{-\bar{k}}|^2 \leq \sup_{\bar{m}} |c_{\bar{m}}|^2 \sum_{\bar{k}} |c_{\bar{k}}|^2 = C_1 < \infty.$$

Therefore as it follows from (28)

$$\sum_{\bar{m}} |a_{\bar{m}}|^2 \leq C_1 \sum_{\bar{m}} \sum_{\bar{k}} |\beta_{\bar{m}}^{\bar{k}}(\varepsilon)|^2. \quad (29)$$

To prove that this double sum tends to zero for $\varepsilon \rightarrow 0$, we use the integral criterion; we denote

$$G(\varepsilon) = \int_{|\bar{k}|>1} dk_1 dk_2 \int_{|\bar{m}+\bar{k}|>1} dm_1 dm_2 |\beta_{\bar{m}}^{\bar{k}}(\varepsilon)|^2.$$

By introducing new variables $s_1 = (\bar{m} + \bar{k}) \times \frac{\bar{k}}{|\bar{k}|}$; $s_2 = (\bar{m} + \bar{k}) \times \frac{\bar{k}^\perp}{|\bar{k}|}$, where $\bar{k}^\perp = (-k_2, k_1)$ we estimate $G(\varepsilon)$ extending integration in s_1, s_2 on the whole $\mathbb{R}^2$

$$G(\varepsilon) \leq \int_{\mathbb{R}^2 \setminus D} dk_1 dk_2 \int_{\mathbb{R}^2} \left(1 - \frac{\sin(\varepsilon |\bar{k}| s_1)}{\varepsilon |\bar{k}| s_1}\right)^2 \frac{s_1^2}{\bar{k}^2} \frac{ds_1 ds_2}{(1 + s_1^2 + s_2^2)^\sigma (1 + \bar{k}^2)^\sigma},$$

where D is the unit disc on $\mathbb{R}^2$, $\bar{k}^2 < 1$. For $\sigma > 2$ the integral over s_2 is finite and we have

$$\int_{\mathbb{R}} \frac{ds_2}{(1 + s_1^2 + s_2^2)^\sigma} = \frac{M_\sigma}{(1 + s_1^2)^{\sigma-1/2}},$$

where the constant M_σ grows to infinity for $\sigma \rightarrow 2$. Therefore introducing $z = \varepsilon |\bar{k}| s_1$ we have

$$G(\varepsilon) \leq \frac{M_\sigma}{\varepsilon^3} \int_{\mathbb{R}^2 \setminus D} \frac{dk_1 dk_2}{\bar{k}^2 (1 + \bar{k}^2)^\sigma} \int_{\mathbb{R}^1} \left(1 - \frac{\sin z}{z}\right)^2 \frac{z^2 dz}{(1 + \frac{z^2}{\varepsilon^2 \bar{k}^2})^{\sigma-1/2}}.$$

Noting that $|1 - \frac{\sin z}{z}| \leq \min(\frac{z^2}{6}, 2)$ we can estimate the integral over dz splitting it in two parts as follows

$$\begin{aligned} \int_0^\infty (1 - \frac{\sin z}{z})^2 \frac{z^2 dz}{(1 + \frac{z^2}{\varepsilon^2 k^2})^{\sigma-1/2}} &\leq \frac{1}{36} \int_0^{|\bar{k}\varepsilon|^\alpha} z^6 dz + \left| 4 \int_{|\bar{k}\varepsilon|^\alpha}^\infty |\varepsilon \bar{k}|^{2\sigma-1} z^{3-2\sigma} dz \right| = \\ &= \frac{1}{252} |\bar{k}\varepsilon|^{7\alpha} + \frac{2}{|2-\sigma|} |\varepsilon \bar{k}|^{2\sigma-1+2\alpha(2-\sigma)}. \end{aligned}$$

Taking $\alpha = \frac{2\sigma-1}{2\sigma+3}$ we make both exponents equal to $7\frac{2\sigma-1}{2\sigma+3}$ and finally one obtains

$$G(\varepsilon) \leq \varepsilon^r M_\sigma \left\{ \frac{1}{126} + \frac{4}{\sigma-2} \right\} \int_{IR^2 \setminus D} \frac{|\bar{k}|^{7\alpha-4}}{(1+\bar{k}^2)^\sigma} dk^1 dk^2,$$

where $r = 8\frac{\sigma-2}{2\sigma+3}$. For $\sigma > 2$, the integral over k^1, k^2 is finite, and thus $G(\varepsilon) \rightarrow 0$ for $\varepsilon \rightarrow 0$.

In this way we have proved that for $\sigma > 2$ the first term on the r.h.s. of Eq(23) is vanishing when $N \rightarrow \infty$ ($\varepsilon = 2\pi/N$). It converges to zero as $\varepsilon^{r/2}$, i.e. as $N^{-r/2}$, where r depends on σ . Indeed, we have

$$|\sum_{\bar{m}} a_{\bar{m}}(\varepsilon) \Omega_{-\bar{m}}| \leq |\Omega| \left(\sum_{\bar{m}} |a_{\bar{m}}(\varepsilon)|^2 \right)^{1/2} = |\Omega| G(\varepsilon)^{1/2}.$$

Now let us estimate the second term on the r.h.s of (23). Denoting $\bar{\mu} = \bar{m} + \bar{k}$ we have

$$\begin{aligned} \left| \sum_{\bar{m}} \sum_{\bar{k}} \frac{\sin(\varepsilon \bar{m} \times \bar{k})}{\varepsilon \bar{k}^2} \omega_{\bar{m}+\bar{k}} \Omega_{-\bar{k}} \Omega_{-\bar{m}} \right| &\leq \sum_{\bar{\mu}, \bar{k}} \left| \frac{\sin(\varepsilon \bar{\mu} \times \bar{k})}{\varepsilon (\bar{\mu} \times \bar{k})} \frac{\bar{\mu} \times \bar{k}}{\bar{k}^2} \frac{c_{\bar{\mu}}}{(1+\bar{\mu}^2)^{\sigma/2}} \Omega_{-\bar{k}} \Omega_{\bar{k}-\bar{\mu}} \right| \leq \\ &\sum_{\bar{\mu}, \bar{k}} |c_{\bar{\mu}} (1+\bar{\mu}^2)^{\frac{1-\sigma}{2}}| |\Omega_{-\bar{k}} \Omega_{\bar{k}-\bar{\mu}}| \leq K_\sigma \|\Omega\|^2, \end{aligned}$$

where $K_\sigma = \sum_{\bar{\mu}} |c_{\bar{\mu}} (1+\bar{\mu}^2)^{\frac{1-\sigma}{2}}|$. From Schwartz inequality K_σ is finite for $\sigma > 2$.

Now let us come back to Eq.(23). Dividing both sides by $\|\Omega(t)\|$ and integrating with respect to time we come to

$$\begin{aligned} \|\Omega(t)\| &= \|\Omega(0)\| + \int_0^t \frac{dt}{\|\Omega(t)\|} \sum_{\bar{m}} a_{\bar{m}}(\varepsilon) \Omega_{\bar{m}}(t) + \\ &\int_0^t \frac{dt}{\|\Omega(t)\|} \sum_{\bar{m}, \bar{k}} \frac{\sin(\varepsilon \bar{m} \times \bar{k})}{\varepsilon \bar{k}^2} \omega_{\bar{m}+\bar{k}} \Omega_{-\bar{m}} \Omega_{-\bar{k}}. \end{aligned}$$

Applying the estimations which we had derived one arrives at

$$\| \Omega(t) \| \leq \left\{ \| \Omega(0) \| + t G^{1/2}(\varepsilon) \right\} + \int_0^t K_\sigma \| \Omega(t) \| dt,$$

which by Gronwall inequality implies that on the time interval $(0, T)$ we have

$$\| \Omega(t) \| \leq \left\{ \| \Omega(0) \| + T G^{1/2}(\varepsilon) \right\} e^{K_\sigma t}, \quad 0 \leq t \leq T.$$

For $N \rightarrow \infty$ the difference $\Omega(0)$ between $\omega(0)$ and $\tilde{\omega}(0)$ tends to zero, at least as $\sim N^{-(\sigma-1)}$, $\| \Omega(0) \| \sim N^{-(\sigma-1)} \rightarrow 0$, similarly, $G(\varepsilon) \sim N^{-r} \rightarrow 0$, and therefore also $\| \Omega(t) \| \rightarrow 0$, which proves that the solution $\tilde{\omega}$ of Euler equation on $su(N)$ approaches the corresponding hydrodynamic solution ω .

In a similar way one can prove a little more general theorem which summarizes our considerations.

Theorem If $\omega(t) \in C^1([0, T], H^\sigma(T^2))$ is a solution of the Euler equation of hydrodynamics on the torus T^2 then corresponding solutions $\tilde{\omega}^N$ (i.e. such that $\tilde{\omega}_m^N(0) = \omega_m^N(0)$ for $-\frac{N-1}{2} \leq m_i < \frac{N-1}{2}$) of the $su(N)$ Euler equations are approaching ω in the norm of $H^k(T^2)$ for every k satisfying $0 \leq k < \sigma - 2$.

4 Numerical simulations

The theorem proved in the last section shows that for large N the difference between solutions of the Euler equation on $su(N)$ and the Galerkin approximation of the Euler equation for ideal incompressible fluid is negligibly small. Numerical schemes usually applied to solve Euler equation are based on the Galerkin approximation. Numerical calculations were performed to test if the group-theoretic approximation could be applied to create a better numerical scheme for solving the Euler equation. There seems to be one important advantage of the group-theoretic finite-mode approach over the Galerkin one. Namely, the former possesses a big number of conserved quantities, which grows with N . The Galerkin approximation does not have this property. As a

result of this difference one can expect higher stability of the group-theoretic approximation for long times.

In the computer simulations we adopted similar initial conditions as Dowker and Wolski [2]. This means we took $\omega(\bar{x}, t = 0) \sim N^2 \delta_{\bar{x}, \bar{x}_0} - 1$. Strictly speaking, such initial conditions are not satisfying the assumptions of our theorem. However, in both approximations, it means the Galerkin and the group-theoretic ones, these initial conditions lead to numerical solutions quite well approximated by the discrete ones. Dowker and Wolski [2] were performing their simulations at the level of stream functions, we were doing that at the level of vorticities.

Calculations in [2] were done on a Hewlett-Packard workstation. We agree with [2] that it is very difficult to go to N s big enough for a convincing numerical illustration of the convergence proved in the theorem. The complexity of the system of ODEs strongly depends on N . In our case the number of equations grows linearly with N , the number of quadratic terms in every equation grows with N^2 . The N is limited not only by the execution time, but also, and in fact mainly, by the memory requirements. This is why we hope that our method will be much more effective in the future — we witness every year significant progress in memory availability.

It was not our aim at this point to make calculations for the biggest possible N , but rather to observe the numerical stability of both methods. This is why calculations were done on a PC (Pentium II, 700MHz, 1GB RAM), in Mathematica 4.1. We restricted ourselves to small N s (relatively easy simulations). The effect is sufficiently strong to be observed at $N = 11, 15, 25$. It is expected from theoretical considerations to be much more significant for big N .

We observed individual Fourier modes as functions of time for both approximations. We were testing time after which the symmetry $\omega_k^* = \omega_{-\bar{k}}$,

(for our choice of initial conditions (real $\omega_{\bar{k}}$) it was $\omega_{\bar{k}} = \omega_{-\bar{k}}$) was breaking. One can easily observe big difference in times of destabilization between the two approximations. For $N = 11$ and $N = 15$ numerical calculations based on the Galerkin method destabilize after the amount of time approximately equal to 4.5 computer units, while the group-theoretic method presented stable behavior till 20 computer time units. For $N = 25$ the difference was significantly bigger. Namely, the group-theoretic method destabilized after about 18 units of time, while the Galerkin method destabilized after 1.5 units of time. In this case the group-theoretic method was more stable than the Galerkin one by a factor of 12. This difference was significant, and we are convinced that with bigger N and longer simulation times the difference would increase. We used in both cases exactly the same initial conditions, numerical methods, numerical accuracy and all other parameters.

As an example, the results of the numerical simulations for the representative cases of $N = 11, 15, 25$ for a pair of modes each in the Galerkin and in the group-theoretic approximations are shown in Figs.1-3. In the graphs is presented the time dependence of the amplitudes of three particular pairs of modes with the numbers $\bar{k}$ and $-\bar{k}$. They are representative for all the pairs of modes. We were looking for the instant of time when the two modes begin to behave differently showing that reliability of the simulations is lost.

5 Conclusions

In this paper we studied the approximate ideal fluid description based on the observation concerning the convergence of the structure constants of the $su(N)$ (N odd) algebras to the structure constants of the algebra $sdiff(T^2)$, and on similarity of the Euler equations defined on the dual algebras of $su(N)$ and $sdiff(T^2)$, respectively. The hypothesis stated in the literature that the Euler equations for $su(N)$ converge to those for $sdiff(T^2)$ was carefully ex-

amined.

The results are:

In addition to the fact known from the literature [13] that the algebras $su(\infty)$ and $sdiff(T^2)$ are not isomorphic, we pointed another difficulty of the approximation. Namely, we have shown that in the limit $N \rightarrow \infty$ renormalized elements of the basis of $su(N)$, used to demonstrate similarity of the structure constants of $\lim_{N \rightarrow \infty} su(N)$ and of $sdiff(T^2)$, become elements of infinite norm.

We proved the statement about the convergence of the solutions of the Euler equations for $su(N)$ to the solutions of the Euler equations for $sdiff(T^2)$. This way the functional analytic part of the approximation eventually found a complete proof.

We performed extensive numerical simulations based on the group-theoretic and the Galerkin finite-mode approximations. They showed that the group-theoretic method is significantly more reliable for long evolution times than the Galerkin one. When the same numerical conditions are applied, the Galerkin method leads to instabilities after evolution time at least a few times shorter than the group-theoretic one. This difference grows with the number of modes used in simulation. For $N=25$ group-theoretic numerical solutions were 12 times more stable than the Galerkin ones. We expect the difference to be sharper for bigger numbers of modes involved. On the other hand, the Galerkin method is significantly less machine time and memory consuming than the group-theoretic one. Galerkin method is appropriate for short evolution time calculations, performed with limited computational power. With given machine power at hands one can obtain solutions for short evolution time using the Galerkin approximation for bigger N than using the group-theoretic one.

The long time scale solutions of the Euler equations are used in meteo-

rology, in weather prediction, as well as in studying oceanic flows, to name just a few applications. The group-theoretic method would allow to improve long time scale predictions reliability in the mentioned areas. It currently requires powerful workstations for implementation. With dramatic growth of computational power and the memory size in recent years, in a few years the group-theoretic method should be easier to apply than it is at the moment.

6 Appendix

Noncommutative tori are one of the simplest examples of noncommutative spaces, which are a subject of study in noncommutative geometry. In this appendix we present material on noncommutative tori, which is relevant for the finite-mode approximation discussed in the paper.

The paradigm of noncommutative geometry grew up from the Gelfand-Naimark theorem, accordingly to which the category of Hausdorff, locally compact topological spaces is dual to the category of commutative C^* -algebras. The commutative C^* -algebras are algebras of continuous complex-valued functions (with compact support in the case of noncompact spaces) on the spaces. The C^* -algebras of continuous functions on compact spaces are additionally unital, with the unit given by the constant function equal to one. Constant functions cannot have compact support on noncompact spaces, therefore the corresponding C^* -algebras are nonunital. Thanks to the Gelfand-Naimark theorem, the entire topological information on the spaces is coded in the algebraic properties of commutative C^* -algebras, to which they are dual.

Relaxing the commutativity property on the C^* -algebras locates us then in the field of noncommutative topology. The subject of its study is not some kind of noncommutative spaces, but rather algebras of (continuous) functions on them. These "functions" are actually operators, and the C^* -algebras

are algebras of operators acting in appropriate Hilbert spaces, in general infinite dimensional. This statement is the content of a second theorem by Gelfand and Naimark. (On Gelfand-Naimark theorems one can consult for example the recent book [20]). The fact that coordinates of the virtual noncommutative spaces are not commuting, and the possibility to realize the algebras of "functions" on them as algebras of operators in a Hilbert space, inspired the creators and developers of this area of mathematics to adopt physicists' terminology and to speak about quantum geometry or quantum topology (the latter name in the case of strictly limiting to the C^* -algebras seems more appropriate) and about quantum spaces.

In contrast to topologists, geometers usually prefer to speak about smooth manifolds rather than topological spaces. Therefore, instead of C^* -algebras noncommutative geometry studies $*$ -algebras that are dense in appropriate C^* -algebras exactly in the manner in which algebras of smooth complex valued functions on manifolds are dense in corresponding algebras of continuous functions. The conditions lead to the notion of a pre- C^* -algebra.

The noncommutative torus is a quantum space, which might be described using either C^* -algebraic or $*$ -algebraic language, with appropriate limits in the latter case locating us in the C^* -algebraic framework. Let us shortly review, following [21], [22] the $*$ -algebraic approach to the noncommutative torus. The algebra $\mathcal{A}_\theta$ of smooth "functions" on the noncommutative torus T_θ^2 is the unital $*$ -algebra generated by 2 indeterminates U_1, U_2 with the relation

$$U_1 U_2 = e^{2\pi i \theta} U_2 U_1 \quad (30)$$

The $*$ structure is defined by imposing unitarity on U_1, U_2 , by $U_1^* = U_1^{-1}$, $U_2^* = U_2^{-1}$. A generic element $a \in \mathcal{A}_\theta$ is then of the form

$$a = \sum_{(m,n) \in \mathbb{Z}^2} a_{mn} U_1^m U_2^n \quad (31)$$

, where a_{mn} is a complex valued Schwarz function on $\mathbb{Z}^2$, i.e. a sequence of complex numbers $\{a_{mn} \in \mathcal{C} : (m, n) \in \mathbb{Z}^2\}$, which decreases rapidly at infinity ($\forall k \in \mathbb{N} \quad \|a\|_k := \sup_{m, n \in \mathbb{Z}} |a_{mn}| (1 + |m| + |n|)^k < \infty$).

In this paper we are interested in the case, in which θ is rational, $\theta = M/N$, where M and N are positive integers taken to be relatively prime. Therefore we present further only the rational θ case. In this case $\mathcal{A}_{M/N}$ is the algebra of smooth sections of a twisted matrix bundle over T^2 . Let us briefly describe the construction leading to the twisted bundle following [21]. The bundle is denoted $\mathcal{M}_q \rightarrow T^2$, with $q = e^{2\pi i M/N}$. The fibers of the bundle are isomorphic with the algebra $Mat(N, \mathcal{C})$ of all $N \times N$ complex-valued matrices. Its transition functions have values in $GL(N, \mathcal{C})$. They act on $Mat(N, \mathcal{C})$ by conjugation.

First, we consider a trivial bundle (we identify here and below T^2 with $S^1 \times S^1$)

$$S^1 \times S^1 \times Mat(N, \mathcal{C}) \xrightarrow{pr_{1,2}} S^1 \times S^1 \quad (32)$$

where $pr_{1,2}$ is the natural projection on the first two members. The algebra of smooth sections of this bundle is

$$C^\infty(S^1 \times S^1, Mat(N, \mathcal{C})) = C^\infty(S^1 \times S^1, \mathcal{C}) \otimes Mat(N, \mathcal{C}) \quad (33)$$

The algebra $Mat(N, \mathcal{C})$ may be understood as the unique algebra generated by 2 unitary elements U_0 and V_0 such that

$$U_0 V_0 = q V_0 U_0, \quad U_0^N = V_0^N = \mathbb{1} \quad (34)$$

The matrices U_0 and V_0 may be represented in the form, which is very closely related to the matrices g and h introduced in the section 2. Here however, the parameter q differs from the parameter λ , since $q = e^{2\pi i M/N}$, but $\lambda = e^{4\pi i/N}$. The algebra of smooth sections of the bundle (32) is generated by unitary commuting elements u , v , and by unitaries U_0 , V_0 generating $GL(N, \mathcal{C})$.

In order to define the twisting of this bundle, the following actions should be considered: the action of $\mathbb{Z}_N \times \mathbb{Z}_N = \mathbb{Z}_N^2$ on $S^1 \times S^1$, $(u, v) \mapsto (q^m u, q^n v)$, $m, n \in \mathbb{Z}_N$, and the action of $\mathbb{Z}_N^2$ on $Mat(N, \mathcal{C})$, $A \mapsto U_0^n V_0^{-m} A V_0^m U_0^{-n}$, $m, n \in \mathbb{Z}_N$.

Then, we impose the condition that the following diagram is commutative

$$\begin{array}{ccc} S^1 \times S^1 \times Mat(N, \mathcal{C}) & \xrightarrow{\mathbb{Z}_N^2} & \mathcal{M}_q \\ pr_{1,2} \downarrow & & \downarrow p_q \\ S^1 \times S^1 & \xrightarrow{\mathbb{Z}_N^2} & S^1 \times S^1 \end{array} \quad (35)$$

There is a vector bundle over $S^1 \times S^1$, $\mathcal{E}_q \rightarrow S^1 \times S^1$ such that $\mathcal{M}_q = End(\mathcal{E}_q)$. This fact is closely related to the Morita equivalence of the non-commutative torus with the commutative one. Namely, Morita equivalence of two C^* -algebras or two pre- C^* -algebras means that the algebras have a common module, on which one of them acts from the left and the other from the right. This is exactly the case for a noncommutative torus with rational "deformation parameter" and the standard commutative torus. We may treat the space of (smooth) sections of the bundle $\mathcal{E}_q \rightarrow S^1 \times S^1$ as the left module for the algebra $\mathcal{M}_q$, since the latter consists of endomorphisms of the former. Simultaneously, since $\mathcal{E}_q \rightarrow S^1 \times S^1$ is a bundle over the standard torus, the pre- C^* -algebra of smooth functions on the torus acts on the space of smooth sections of $\mathcal{E}_q$ from the right. This is the way in which the rational noncommutative tori are related via Morita equivalence with the commutative torus.

However, the Morita equivalence of the two kind of spaces, although very important, is not directly responsible for the convergence effects we studied in the body of this paper. In the geometric picture the limiting procedure may be described in the following way: The algebra $Mat(N, \mathcal{C})$, which is isomorphic with the fibers of the bundle $\mathcal{M}_q \rightarrow S^1 \times S^1$ may be understood as the Lie algebra $gl(N, \mathcal{C})$. From this moment on one should just remind the construction described in the body of the paper.

7 Acknowledgements

This research reported in this paper was begun and to high extend developed when two of the authors (HM and RO) were working at the group T-13 of the Theoretical Division of the Los Alamos National Laboratory, and finished when the two authors were working at the Environmental Science and Waste Technology Division of the Laboratory. The work of Z.P. was supported by Polish State Committee for Scientific Research (KBN) grant *N^o 2PO3B 169 08 1109/PO3/95/08*. Z.P. acknowledges also hospitality of Los Alamos National Laboratory during his visit in October '97. All the authors would like to express their deep gratitude to Donna Spitzmiller, for her warm hospitality during their stay at Theoretical Division.

The work of R.O. was partially supported by a fellowship from the Fulbright Foundation.

H.M. was partially supported by a Directors' Funded Postdoctoral Fellowship offered by Los Alamos National Laboratory.

The authors are grateful to I.Szczyrba for very helpful discussions of issues concerning infinite dimensional Lie algebras and to K. Piechór for long discussions concerning details of the proof.

The authors express their gratitude to S.Shkoller, who turned their attention to a recent paper [19], which was written after their work was completed, and which contains results on the functional analytic convergence similar to ours.

References

- [1] V. Zeitlin, *Finite-mode analogs of 2D ideal hydrodynamics: Coadjoint orbits and local canonical structure*, Physica **D 49**, (1991), 353-362.

- [2] J. S. Dowker, A. Wolski, *Finite model of two-dimensional ideal hydrodynamics*, Phys.Rev., **A 46**, (1992), 6417-6430.
- [3] A. Rouhi, H.D.I. Abarbanel *Symmetric truncations of the shallow-water equations*, Phys.Rev., **E 48**, (1993), 3643-3655.
- [4] R.I. McLachlan, I. Szunyogh, V. Zeitlin, *Hamiltonian finite-dimensional models of baroclinic instability*, Phys.Lett., **A 229**, (1997), 299-305.
- [5] J.Hoppe,
Diffeomorphism Groups, Quantization, and $SU(\infty)$, Int.J.Mod.Phys., **A 4**, (1989), 5235-5248.
- [6] D.B. Fairlie, P. Fletcher, C.K. Zachos, *Trigonometric Structure Constants for New Infinite-Dimensional Algebras*, Phys.Lett., **B 218**, (1989), 203-206.
- [7] E.G. Floratos, J. Iliopoulos, *A Note on the Classical Symmetries of the Closed Bosonic Membranes*, Phys.Lett., **B 201**, (1988), 237-240.
- [8] D.B. Fairlie, C.K. Zachos, *Infinite Dimensional Algebras, Sine Brackets, and $SU(\infty)$* , Phys.Lett., **B 224**, (1989), 101-107.
- [9] P.J. Channel, J.C. Scovel, *Integrators for Lie-Poisson dynamical systems*, Physica, **D 50**, (1991), 80-88.
- [10] R.I. McLachlan, *Explicit Lie-Poisson Integration and the Euler Equations*, Phys. Rev. Lett., **71**, (1993), 3043-3046.
- [11] C. Scovel, A. Weinstein, *Finite Dimensional Lie-Poisson Approximations to Vlasov-Poisson Equations*, Comm. Pure Appl. Math., **47**, (1994), 683-709.
- [12] Z. Ge, C. Scovel, *Hamiltonian Truncation of the Shallow Water Equation*, Lett. Math. Phys., **31**, (1994), 1-13.

- [13] M. Bordemann, J. Hoppe, P. Schaller, M. Schlichenmaier, *$gl(\infty)$ and Geometric Quantization*, Commun. Math. Phys **138**, (1991), 209-244.
- [14] C.N. Pope, K.S. Stelle, *$SU(\infty)$, $SU_+(\infty)$ and Area-Preserving Algebras*, Phys.Lett., **B 226**, (1989), 257-263.
- [15] G.A. Goldin, R. Menikoff, D.H. Sharp, *Diffeomorphism Groups and Quantum Vortex Filaments*, Phys. Rev. Lett., **58**, (1987), 2162-2164.
- [16] G.A. Goldin, R. Owczarek, D.H. Sharp, *Quantum Kinematics of Bosonic Vortex Loops*, in preparation.
- [17] V.I. Arnold, *Mathematical Methods of Classical Mechanics*, Springer-Verlag, Berlin (1978).
- [18] P.-L. Lions, *Mathematical Topics in Fluid Mechanics*, vol.1, Clarendon Press, Oxford (1996).
- [19] I. Gallagher, *Mathematical Analysis of a Structure-Preserving Approximation of the Bidimensional Vorticity Equation*, preprint
- [20] J.M. Gracia-Bondia, J.C. Varilly, H. Figueroa, *Elements of Noncommutative Geometry*, Birkhaeuser Advanced Texts, (2001)
- [21] G. Landi, F. Lizzi, R.J. Szabo, *From Large N Matrices to the Noncommutative Torus*, Comm. Math. Phys., **217**, (2001), 181-201
- [22] M. Dubois-Violette, A. Kriegl, Y. Maeda, P.W. Michor, *Smooth *-Algebras*, preprint, arXiv: math.QA/0106150, (2001)

Figure captions

Fig 1.1 Group - theoretic method, $N=11$, the modes compared:

- i) mode $(-5, 0)$ and mode $(5, 0)$
- ii) mode $(-3, 2)$ and mode $(3, -2)$
- iii) mode $(-3, 0)$ and mode $(3, 0)$

The simulation was still stable at 20 units of time. After this time the $\mathbf{k}$ and $-\mathbf{k}$ modes started to differ from each other, which means that the solution is no longer stable at this point.

Fig 1.2 Galerkin method, $N=11$, the modes compared:

- i mode $(-5, 0)$ and mode $(5, 0)$
- ii mode $(-3, 2)$ and mode $(3, -2)$
- iii mode $(-3, 0)$ and mode $(3, 0)$

The simulation was stable till 4.8 units of time. After this time the $\mathbf{k}$ and $-\mathbf{k}$ modes started to differ from each other, which means that the solution is no longer stable at this point.

Fig 2.1 Group - theoretic method, $N=15$, the modes compared:

- i) mode $(-7, 0)$ and mode $(7, 0)$
- ii) mode $(-1, 2)$ and mode $(1, -2)$
- iii) mode $(-7, 5)$ and mode $(7, -5)$

The simulation was stable till 23 units of time. After this time the $\mathbf{k}$ and $-\mathbf{k}$ modes started to differ from each other, which means that the solution is no longer stable at this point.

Fig 2.2 Galerkin method, $N=15$, the modes compared:

- i) mode $(-7, 0)$ and mode $(7, 0)$
- ii) mode $(-1, 2)$ and mode $(1, -2)$
- iii) mode $(-7, 5)$ and mode $(7, -5)$

The simulation was stable till 2.7 units of time.


Fig 3.1 Group-theoretic method, $N=25$, the modes compared:

- i) mode $(-12, 2)$ and mode $(12, -2)$
- ii) mode $(-6, 2)$ and mode $(6, -2)$
- iii) mode $(-5, 2)$ and mode $(5, -2)$ The simulation was stable till 18 units of time. After this time the $\mathbf{k}$ and $-\mathbf{k}$ modes started to differ from each other, which means that the solution is no longer stable at this point.


Fig 3.2 Galerkin method, $N=25$, the modes compared:

- i) mode $(-12, 2)$ and mode $(12, -2)$
- ii) mode $(-6, 2)$ and mode $(6, -2)$
- iii) mode $(-5, 2)$ and mode $(5, -2)$


The simulation was stable till 1.5 units of time. After this time the $\mathbf{k}$ and $-\mathbf{k}$ modes started to differ from each other, which means that the solution is no longer stable at this point.




$N=11$, modes $(-5,0), (5,0)$




$N=11$, modes $(-3,2), (3,-2)$




$N=11$, modes $(-3,0), (3,0)$




$N=11$, modes $(-5,0), (5,0)$, Galerkin




$N=11$, modes $(-3,2)$, $(3,-2)$, Galerkin




$N=11$, modes $(-3,0), (3,0)$, Galerkin




$$N=15, \text{ modes } (-7,0), (7,0)$$




$$N=15, \text{ modes } (-1,2), (1,-2)$$




$N=15$, modes $(-7,5), (7,-5)$




$N=15$, modes $(7,0), (-7,0)$, Galerkin




$N=15$, modes $(-1,2), (1,-2)$, Galerkin




$N=15$, modes $(7,-5), (-7,5)$, Galerkin




$N=25$, modes $(-12, 2), (12, -2)$




$N=25$, modes $(-6,2), (6,-2)$, Galerkin




$N=25$, modes $(-5, 2)$, $(5, -2)$



$N=25$, modes $(-12, 2), (12, -2)$, Galerkin



$$N=28, \text{ modes } (-6, 2), (6, -2)$$



$N=25$, modes $(-5,2), (5,2)$, Galerkin